

Electron-correlation effects in the g -factor of light Li-like ions

V. A. Yerokhin,¹ K. Pachucki,² M. Puchalski,^{2,3} Z. Harman,⁴ and C. H. Keitel⁴

¹*Center for Advanced Studies, Peter the Great St. Petersburg Polytechnic University, Polytekhnicheskaya 29, 195251 St. Petersburg, Russia*

²*Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

³*Faculty of Chemistry, Adam Mickiewicz University, Umultowska 89b, 61-614 Poznań, Poland*

⁴*Max Planck Institute for Nuclear Physics, Saupfercheckweg 1, D 69117 Heidelberg, Germany*

We investigate electron-correlation effects in the g -factor of the ground state of Li-like ions. Our calculations are performed within the nonrelativistic quantum electrodynamics (NRQED) expansion up to two leading orders in the fine-structure constant α , α^2 and α^3 . The dependence of the NRQED results on the nuclear charge number Z is studied and the individual $1/Z$ -expansion contributions are identified. Combining the obtained data with the results of the all-order (in $Z\alpha$) calculations performed within the $1/Z$ expansion, we derive the unified theoretical predictions for the g -factor of light Li-like ions.

INTRODUCTION

Measurements of the bound-electron g -factor in light H-like ions [1–3] provide one of the best tests of the bound-state QED theory as well as the most accurate determination of the electron mass [4]. Similar experiments on Li-like ions [5, 6] probe the QED theory of the electron-correlation effects. In future, a combination of g -factor experiments on Li-like and H-like ions can be used as a new way to determine the fine-structure constant α [7].

The QED effects in the g -factors of few-electron atoms can be systematically treated within the two methods. The first method starts with the Dirac equation for the valence electron in a Coulomb field of the nucleus and accounts for the radiative and electron-electron interaction effects by perturbation theory. The expansion parameter for the electron-electron interaction is $1/Z$ (where Z is the nuclear charge number). This method accounts for all orders in the nuclear binding strength parameter $Z\alpha$ and thus is most effective for high- Z atoms. Extensive QED calculations of the g -factors of Li-like ions within the $1/Z$ expansion method were performed by Shabaev and co-workers [8–12].

The starting point of the second method is the Schrödinger equation that includes both the electron-nucleus and the electron-electron Coulomb interactions. The radiative and relativistic effects are accounted for by perturbation theory, with the expansion parameters α and $Z\alpha$, respectively. This method is often denoted as the Nonrelativistic Quantum Electrodynamics (NRQED) approach, since the coefficients of the perturbation expansion can be derived systematically within NRQED. In contrast to the first method, the NRQED treatment accounts for all orders in $1/Z$ but expands in $Z\alpha$ and thus is most effective for low- Z atoms. Calculations of g -factors by this method were carried out by Hegstrom [13] and, more recently, by Yan [14, 15].

The experiments on the g -factor of Li-like atoms have been so far performed in the intermediate region of Z , where the two methods are complementary to each other.

The optimal theoretical treatment in this region of Z can be achieved by combining them together. To this end, one would need to identify (i) the individual $1/Z$ -expansion terms in the NRQED calculations and (ii) the individual $Z\alpha$ -expansion terms in the $1/Z$ -expansion results. A combination of these results would then provide a unified theory. The goal of the present investigation is to make the first steps along this path.

In this work we perform the NRQED calculation of the leading relativistic ($\sim \alpha^2$) and the leading QED ($\sim \alpha^3$) corrections to the g -factor of the ground state of Li-like ions, extending previous calculations by Yan to a larger region of Z and improving the numerical accuracy. We identify the individual $1/Z$ expansion terms of these corrections. In particular, we obtain the higher-order electron-correlation contribution of the relative order $1/Z^3$ and higher, thus removing one of the dominant sources of the uncertainty of theoretical predictions [12].

I. NRQED APPROACH

Within the NRQED approach, the bound-electron g -factor of a light atom is represented as an expansion in powers of the fine-structure constant α ,

$$g = g_e + \alpha^2 g^{(2)} + \alpha^3 g^{(3)} + \alpha^4 g^{(4)} + \dots, \quad (1)$$

where g_e is the free-electron g -factor and $g^{(n)}$ are the binding corrections. The expansion coefficients $g^{(n)}$ can be further expanded in powers of the electron-to-proton mass ratio m/M ,

$$g^{(n)} = g_\infty^{(n)} + \frac{m}{M} g_M^{(n)} + \dots \quad (2)$$

The interaction of a *free* nonrelativistic electron with a constant external magnetic field \vec{B} is described by the Hamiltonian

$$\begin{aligned} H &= \mu_B (1 + \kappa) \vec{\sigma} \cdot \vec{B} \\ &= \mu_B (1 + \kappa) 2 \vec{s} \cdot \vec{B} \equiv \mu_B g_e \vec{s} \cdot \vec{B}, \end{aligned} \quad (3)$$

where $\mu_B = -e/(2m)$ is the Bohr magneton, $\vec{\sigma}$ is the vector of Pauli matrices, \vec{s} is the electron spin operator, κ is the anomalous magnetic moment of the free electron, which is connected to the free-electron g -factor by $g_e \equiv 2(1 + \kappa) = 2 + \alpha/\pi + \dots$

Many years ago Hegstrom [13] derived the Hamiltonian describing the interaction of an atom with the magnetic field, which accounts for the leading relativistic, QED, and nuclear recoil effects. The resulting Hamiltonian is complete through orders of α^2 , α^3 , $\alpha^2 m/M$, and $\alpha^3 m/M$. The corresponding numerical calculations for Li-like atoms were performed by Yan [14, 15].

In the present work, we address the leading relativistic and QED corrections to the g -factor of Li-like atoms.

These corrections are induced by the effective Hamiltonian δH , which, for the case of the S states, can be simplified to take a very compact form,

$$\delta H = \sum_a \mu_B Q_a \vec{\sigma}_a \cdot \vec{B}, \quad (4)$$

$$Q_a = Q_a^{(2)} + \kappa Q_a^{(3)}, \quad (5)$$

$$Q_a^{(2)} = \frac{1}{3} \left(-2p_a^2 + \frac{Z}{r_a} - \sum_{b \neq a} \frac{1}{r_{ab}} \right), \quad (6)$$

$$Q_a^{(3)} = \frac{1}{3} \left(-\frac{p_a^2}{2} + \frac{Z}{r_a} - \sum_{b \neq a} \frac{1}{r_{ab}} \right), \quad (7)$$

where the indices a and $b = (1, 2, 3)$ numerate the electrons in the atom.

Expectation values of the operators Q_a are evaluated with the nonrelativistic wave function ψ . This function is the antisymmetrized product (\mathcal{A}) of the spacial function ϕ and the spin function χ ,

$$\psi = \mathcal{A}[\phi(\vec{r}_1, \vec{r}_2, \vec{r}_3) \chi], \quad (8)$$

$$\chi = [\alpha(1) \beta(2) - \beta(1) \alpha(2)] \alpha(3), \quad (9)$$

where $\sigma_z \alpha(\cdot) = \alpha(\cdot)$ and $\sigma_z \beta(\cdot) = -\beta(\cdot)$. Matrix elements of a spin-independent operator H , after eliminating spin variables, can be expressed as

$$\langle \psi' | H | \psi \rangle = \left\langle \phi'(r_1, r_2, r_3) \left| H \left[2\phi(r_1, r_2, r_3) + 2\phi(r_2, r_1, r_3) - \phi(r_2, r_3, r_1) - \phi(r_3, r_2, r_1) - \phi(r_3, r_1, r_2) - \phi(r_1, r_3, r_2) \right] \right. \right\rangle. \quad (10)$$

Matrix elements of the spin-dependent operators are expressed as

$$\langle \psi' | \sum_a Q_a \vec{\sigma}_a | \psi \rangle = \sum_a \langle \phi' | Q_a | \phi \rangle_F 2\vec{S}, \quad (11)$$

where $\vec{S} = \sum_a \vec{s}_a$ and

$$\begin{aligned} \sum_a \langle \phi' | Q_a | \phi \rangle_F &= \left\langle \phi'(r_1, r_2, r_3) \left| 2Q_3 [\phi(r_1, r_2, r_3) + \phi(r_2, r_1, r_3)] - (Q_1 - Q_2 + Q_3) [\phi(r_2, r_3, r_1) + \phi(r_3, r_2, r_1)] \right. \right. \\ &\quad \left. \left. - (Q_2 - Q_1 + Q_3) [\phi(r_1, r_3, r_2) + \phi(r_3, r_1, r_2)] \right. \right\rangle. \end{aligned} \quad (12)$$

The corresponding corrections to the g -factor are

$$g_\infty^{(2)} = 2 \sum_a \langle Q_a^{(2)} \rangle_F, \quad (13)$$

$$g_\infty^{(3)} = \frac{1}{\pi} \sum_a \langle Q_a^{(3)} \rangle_F. \quad (14)$$

We calculated the matrix elements (13) and (14) by using accurate variational wave functions in the Hyleraas basis. The method is described in our previous investigations [16–18]. Our numerical results for $g_\infty^{(2)}$ and $g_\infty^{(3)}$ are presented in Tables I and II, respectively. The values listed in the tables were obtained by using the basis with the expansion parameter $\Omega = n_1 + n_2 + n_3 + n_4 + n_5 + n_6 = 12$. The specified uncertainties were obtained by taking

the differences of the results with $\Omega = 12$ and 11. For lithium, we find good agreement with the previous calculations by Yan [14, 15], our results being several digits more accurate. For lithium-like ions we observe small deviations outside of the Yan's error bars.

A. Relativistic correction $g_\infty^{(2)}$

The leading relativistic correction of order α^2 can be expanded in $1/Z$ as follows

$$g_\infty^{(2)}(Z) = -\frac{Z^2}{6} + \frac{940}{2187} Z + c^{(2,0)} + \frac{H^{(2,-1)}(Z)}{Z}, \quad (15)$$

where $H^{(2,-1)}(Z)$ is the remainder function that incorporates all higher orders in $1/Z$, $H^{(2,-1)}(Z) \rightarrow c^{(2,-1)}$ as $Z \rightarrow \infty$. The leading coefficient of the expansion (15) follows from the hydrogenic limit summarized in Appendix A, whereas the second coefficient is derived in the present work. The higher-order coefficients were obtained by fitting the numerical results for $g_\infty^{(2)}$, as described in Appendix B.

Our fitting results for the first higher-order expansion coefficients are

$$c^{(2,0)} = -0.128\,204(9), \quad c^{(2,-1)} = 0.028\,78(46). \quad (16)$$

We would like to stress that in order to achieve such precision of the fitted coefficients, it was important to have highly accurate numerical results for $g_\infty^{(2)}$ for a sufficiently wide range of Z . In particular, if we apply the same fitting procedure to the analogous results of Yan [14, 15], we get results consistent with Eq. (16) but much less accurate.

Using the result for $c^{(2,0)}$ from Eq. (16), we can extract the remainder function $H^{(2,-1)}(Z)$ from our numerical data for $g_\infty^{(2)}$. The corresponding results are presented in the last column of Table I. The errors of the listed values of $H^{(2,-1)}$ come from the uncertainty of $c^{(2,0)}$. In the case of silicon, we obtain $H^{(2,-1)}(14) = 0.024\,77(13)$, which agrees with but is more precise than the corresponding result of 0.0244(15) obtained by the Configuration-Interaction Dirac-Fock (CI-DF) method in Ref. [12]. We note that $H^{(2,-1)}$ previously yielded one of the two main errors of the total theoretical g -factor predictions.

B. QED correction $g_\infty^{(3)}$

The leading QED correction of order α^3 can be expanded in $1/Z$ as follows

$$g_\infty^{(3)}(Z) = \frac{1}{24\pi} Z^2 - \frac{274}{2187\pi} Z + H^{(3,0)}(Z), \quad (17)$$

where $H^{(3,0)}(Z)$ is the remainder that incorporates the higher orders in $1/Z$, $H^{(3,0)} \rightarrow c^{(3,0)}$ as $Z \rightarrow \infty$. The leading coefficient of the expansion (17) comes from the hydrogenic limit, Eq. (A2), whereas the second term was derived in Ref. [9].

Using the known results for the first two terms of the expansion (17), we identify values of the remainder function $H^{(3,0)}(Z)$ from our numerical results for $g_\infty^{(3)}$, with the corresponding results presented in the last column of Table II. In particular, for silicon we obtain $H^{(3,0)}(14) = 0.022\,4679$, which agrees with the corresponding value of 0.0224(10), obtained in Ref. [9] by fitting the results of Yan [14, 15].

Our fitting results for the first expansion coefficients of $H^{(3,0)}(Z)$ are

$$c^{(3,0)} = 0.022\,412(2), \quad c^{(3,-1)} = 0.000\,53(7). \quad (18)$$

These results can be used for estimating the $H^{(3,0)}(Z)$ function for higher values of Z .

We note that $g_\infty^{(3)}$ is induced by the one-loop part of the anomalous magnetic moment (AMM), α/π . According to Eq. (5), analogous corrections due to the n -loop part of the AMM differ from $g_\infty^{(3)}$ only by a prefactor. In particular, the two-loop part of $g_\infty^{(4)}$ is

$$g_{\text{two-loop}}^{(4)} = \frac{2A_2}{\pi} g_\infty^{(3)}, \quad (19)$$

where A_2 is the two-loop contribution to the AMM defined in Eq. (A6).

C. Recoil correction $g_M^{(2)}$

The leading recoil correction of order $\alpha^2 m/M$ can be expanded in $1/Z$ as follows

$$g_M^{(2)}(Z) = \frac{1}{4} Z^2 + Z H_M^{(2,1)}(Z), \quad (20)$$

where the leading coefficient follows from the hydrogenic limit, Eq. (A8), and $H_M^{(2,1)}(Z)$ is the higher-order remainder function, $H_M^{(2,1)} \rightarrow c_M^{(2,1)}$ as $Z \rightarrow \infty$.

In the present work we obtain the remainder function and the coefficient $c_M^{(2,1)}$ by fitting the results of Yan [14, 15]. Our value for the coefficient

$$c_M^{(2,1)} = -0.860\,3(8) \quad (21)$$

disagrees with the corresponding result of $-0.825(5)$ from Ref. [9] obtained by fitting the same results of Yan. We do not know the reason for this disagreement. Our fitting procedure was the same as used for the $g_\infty^{(2)}$ and $g_\infty^{(3)}$ corrections and it reproduces well the analytical value of the leading coefficient in Eq. (20). We also obtain the remainder function for silicon as

$$H_M^{(2,1)}(14) = -0.832\,9(1). \quad (22)$$

D. Radiative recoil correction $g_M^{(3)}$

The radiative recoil correction of order $\alpha^3 m/M$ can be expanded in $1/Z$ as follows

$$g_M^{(3)}(Z) = -\frac{1}{12\pi} Z^2 + Z H_M^{(3,1)}(Z), \quad (23)$$

where the leading coefficient follows from the hydrogenic limit, Eq. (A8), and $H_M^{(3,1)}(Z)$ is the higher-order remainder, $H_M^{(3,1)} \rightarrow c_M^{(3,1)}$ as $Z \rightarrow \infty$.

In the present work we obtain the remainder function and the coefficient $c_M^{(3,1)}$ by fitting the results of Yan [14, 15]. Our values for the coefficient and the remainder are

$$c_M^{(3,1)} = 0.040\,23(4), \quad H_M^{(3,1)}(14) = 0.040\,337(6). \quad (24)$$

TABLE I. The leading relativistic contribution $g_\infty^{(2)}$ to the g -factor of the ground state of Li-like atoms and the corresponding higher-order remainder function $H^{(2,-1)}$ defined by Eq. (15).

Z	$g_\infty^{(2)}$	$H^{(2,-1)}$
3	-0.343 332 404 (3) -0.343 332 42 (7) ^a	-0.013 70 (3)
4	-1.074 312 532 (7)	0.005 23 (4)
5	-2.143 265 913 (2)	0.012 71 (5)
6	-3.546 553 940 (2)	0.016 65 (5)
7	-5.283 459 746 (4)	0.019 06 (6)
8	-7.353 784 626 (1)	0.020 69 (7)
9	-9.757 463 309 (3)	0.021 85 (8)
10	-12.494 472 721 (5)	0.022 73 (9)
11	-15.564 804 889 (4)	0.023 4 (1)
12	-18.968 457 638 (1) -18.968 460 5 (2) ^a	0.024 0 (1)
13	-22.705 431 064 (2)	0.024 4 (1)
14	-26.775 726 109 (1)	0.024 8 (1)

^a Ref. [15].

TABLE II. The leading QED contribution $g_\infty^{(3)}$ to the g -factor of the ground state of Li-like atoms and the corresponding higher-order remainder function $H^{(3,0)}$ defined by Eq. (17).

Z	$g_\infty^{(3)}$	$H^{(3,0)}$
3	0.023 071 092 3 (7) 0.023 071 11 (2) ^a	0.023 344
4	0.075 560 527 2 (1)	0.022 873
5	0.154 876 875 2 (2)	0.022 703
6	0.260 805 551 9 (3)	0.022 619
7	0.393 295 230 1 (5)	0.022 570
8	0.552 328 059 8 (1)	0.022 539
9	0.737 896 374 9 (2)	0.022 518
10	0.949 996 389 1 (4)	0.022 502
11	1.188 626 037 9 (1)	0.022 490
12	1.453 784 110 3 (1) 1.453 784 66 (4) ^a	0.022 481
13	1.745 469 851 3 (1)	0.022 474
14	2.063 682 768 1 (1)	0.022 468

^a Ref. [15].

II. RESULTS AND DISCUSSION

The summary of individual binding corrections to the g -factors of Li-like silicon, oxygen, and carbon ions is presented in Tables III, IV and V, respectively. The sum of all binding corrections gives the difference between the g -factor of the atom and the free-electron g -factor, $g - g_e$, which may be compared to the experimental data and to other theoretical predictions by using the experimental value of the free-electron g -factor [19],

$$g_e = 2.002\,319\,304\,361\,(6). \quad (25)$$

In the tables, the columns labelled ‘‘LO’’ present results for the lowest-order (in $Z\alpha$) parts of the corre-

sponding corrections. The columns labelled ‘‘HO’’ contain results for the higher-order remainders, which are suppressed by a factor of $(Z\alpha)^2$ as compared to the corresponding LO part.

The largest contribution to $g - g_e$ comes from the electron-electron interaction. The corresponding LO part is discussed in Sec. IA. The $1/Z^0$ HO term comes from the hydrogenic limit, Eq. (A1). The $1/Z^1$ HO term originates from the one-photon exchange diagrams, first calculated in Ref. [8] and reevaluated in this work to a higher precision. The $1/Z^2$ HO term comes from the two-photon exchange diagrams, which were calculated to all orders in $Z\alpha$ in Ref. [12]. For silicon, we identify the $1/Z^2$ HO term from the all-order numerical result of Ref. [12]. For oxygen and carbon, there were no results reported in there, so we estimate the $1/Z^2$ HO term by scaling the silicon’s result. E.g., for oxygen we obtain

$$\delta g = -0.000\,92\,\alpha^2\,(8/14)^2 = -0.000\,3\,\alpha^2. \quad (26)$$

We ascribe the uncertainty of 50% to this estimation. The $1/Z^3$ HO term is unknown; the corresponding uncertainty was estimated as the $1/Z^3$ LO term multiplied by the ratio of the $1/Z^2$ HO-to-LO terms, and by a conservative factor of 1.5.

The LO part of the one-loop QED correction is discussed in Sec. IB. The corresponding $1/Z^0$ HO term comes from the hydrogenic limit, Eq. (A2). The $1/Z^1$ HO term is induced by the screened QED diagrams, calculated to all orders in $Z\alpha$ in Refs. [11, 12]. For silicon, we take the result presented in Table II of Refs. [12] and identify the $1/Z^1$ contribution by subtracting the $1/Z^{2+}$ part taken from Table V of Ref. [9]. For carbon and oxygen, we scale the silicon result and ascribe a 100% uncertainty to this estimate. The $1/Z^2$ HO term has not been evaluated yet. We estimated its uncertainty as the $1/Z^2$ LO term multiplied by the ratio of the $1/Z^1$ HO and LO terms, and by an additional conservative factor of 1.5.

The LO part of the recoil correction is discussed in Sec. IC. The only HO recoil contribution available today for oxygen and silicon is the $(Z\alpha)^4 m/M$ correction obtained in Ref. [11] in the hydrogenic limit (see Eq. (A8)). We note that a calculation complete to all orders in $Z\alpha$ was reported in Ref. [6], but only for calcium. In the absence of such calculations for other ions, we estimate the uncertainty due to higher orders in $Z\alpha$ on the basis of the results available for the $1s$ state [20].

The $1/Z^0$ part of the two-loop QED correction is given by Eq. (A6), whereas the $1/Z^{1+}$ part is described in Sec. IB. The finite nuclear size correction is taken from our previous investigation [21].

In Tables III, IV and V, we summarize all known theoretical contributions to $g - g_e$ for Li-like silicon, oxygen, and carbon and compare the results with previous theoretical and experimental data. For silicon, we observe a very good agreement with the theoretical prediction by Volotka et al. [12] and with the experimental result [5]. Our prediction is slightly more accurate than that by

Volotka et al., mainly because of the improvement in the $1/Z^{3+}$ electron-correlation correction. For oxygen and carbon, we find a marginal agreement with the previous theoretical calculations of Glazov et al. [9] but improve their accuracy by a factor of 3 (oxygen) or 4 (carbon). The main difference between the results comes from the $1/Z^{2+}$ electron-correlation correction, which was evaluated by the CI-DF method in Ref. [9] and by the NRQED method in the present work.

The largest uncertainty of our theoretical prediction for silicon stems from the $1/Z^1$ part of the one-loop QED effect, also known as the screened QED correction. The corresponding uncertainty is the estimated error of the numerical evaluation [12], which can be improved by dedicated calculations. For oxygen and carbon, the largest theoretical error comes from the $1/Z^2$ part of the electron-electron interaction correction. This error can be eliminated by extending the all-order calculation of the two-photon exchange diagrams by Volotka et al. [12] to lower- Z ions, or by performing the NRQED calculations of the next-order α^4 effect.

Summing up, we have performed NRQED calculations of the electron-correlation effects to the g factor of the ground state of Li-like atoms. By fitting the Z dependence of the NRQED results for the α^2 and α^3 effects and the corresponding recoil corrections, we have identified their individual $1/Z$ -expansion contributions. Combining the obtained data with the results of the all-order (in $Z\alpha$) calculations performed within the $1/Z$ expansion, we have derived unified theoretical predictions for the g -factor of light Li-like ions and improved the theoretical precision.

ACKNOWLEDGEMENT

V.A.Y. acknowledges support by the Ministry of Education and Science of the Russian Federation Grant No. 3.5397.2017/BY. Work of M.P. and K.P. was supported by the National Science Center (Poland) Grant No. 2012/04/A/ST2/00105. Fruitful discussions with D. A. Glazov, V. M. Shabaev, and A. V. Volotka are gratefully acknowledged.

Appendix A: Hydrogenic limit

The g -factor of the ground state of a Li-like atom in the hydrogenic limit (i.e., neglecting the electron-electron interaction) coincides with the g -factor of the $2s$ state of the corresponding H-like ion. In this section we summarize the theory of the g -factor of the hydrogenic $2s$ state.

The relativistic value of the $2s$ g -factor is obtained from the Dirac equation, with the (point-nucleus) result

$$g = \frac{2}{3} \left[1 + \sqrt{2 + 2\sqrt{1 - (Z\alpha)^2}} \right] = 2 - \frac{(Z\alpha)^2}{6} + \dots \quad (\text{A1})$$

The one-loop QED correction (for the point nucleus) is [22–24]

$$g_{\text{QED}}^{(1)} = \frac{\alpha}{\pi} \left\{ 1 + \frac{(Z\alpha)^2}{24} + \frac{(Z\alpha)^4}{8} \left[\frac{32}{9} \ln[(Z\alpha)^{-2}] + b_{40}^{(1)} \right] + \frac{(Z\alpha)^5}{8} H^{(1)}(Z\alpha) \right\}, \quad (\text{A2})$$

where $b_{40}^{(1)} = -11.774\,382\,27$ [23, 24] and $H(Z\alpha)$ is the remainder function that incorporates all higher orders in $Z\alpha$. The self-energy part of the remainder function was obtained numerically in Ref. [25] to be

$$\begin{aligned} H_{\text{SE}}^{(1)}(6\alpha) &= 22.48(1), \\ H_{\text{SE}}^{(1)}(8\alpha) &= 22.221(4), \\ H_{\text{SE}}^{(1)}(14\alpha) &= 21.486(1). \end{aligned} \quad (\text{A3})$$

The vacuum-polarization part of the remainder function consists of the so-called electric-loop and magnetic-loop parts. The electric-loop part is relatively simple and evaluated numerically by many authors, e.g., by us,

$$\begin{aligned} H_{\text{VP,el}}^{(1)}(6\alpha) &= 1.46, \\ H_{\text{VP,el}}^{(1)}(8\alpha) &= 1.388, \\ H_{\text{VP,el}}^{(1)}(14\alpha) &= 1.1996, \end{aligned} \quad (\text{A4})$$

whereas the magnetic-loop part is given by [26, 27]

$$H_{\text{VP,ml}}^{(1)}(Z\alpha) = \frac{7\pi}{216} - (Z\alpha) \frac{8}{135} \left[\ln(Z\alpha) + 2.6 + \frac{5}{8} \right]. \quad (\text{A5})$$

The two-loop QED correction is

$$g_{\text{QED}}^{(2)} = \frac{\alpha^2}{\pi^2} \left\{ 2A_2 + 2A_2 \frac{(Z\alpha)^2}{24} + \frac{(Z\alpha)^4}{8} \left[\frac{28}{9} \ln[(Z\alpha)^{-2}] + b_{40}^{(2)} + \frac{16 - 19\pi^2}{108} \right] \right\}, \quad (\text{A6})$$

where $A_2 = -0.328\,478\,444\,00\dots$ is the two-loop contribution to the electron anomalous magnetic moment and $b_{40}^{(2)} = -17.157\,236\,58$ [23, 24]. The last term $O((Z\alpha)^4)$ in Eq. (A6) is the light-by-light scattering contribution recently calculated in Ref. [28].

The three and higher-loop QED corrections can be summarized as

$$g_{\text{QED}}^{(\geq 3)} = \sum_{n=3}^5 \frac{\alpha^n}{\pi^n} \left\{ 2A_n + 2A_n \frac{(Z\alpha)^2}{24} \right\}, \quad (\text{A7})$$

where [29, 30]

$$\begin{aligned} A_3 &= 1.181\,234\,017\dots, \\ A_4 &= -1.912\,245\,765\dots, \\ A_5 &= 7.79(34). \end{aligned}$$

The recoil correction, including the second-order recoil $O((m/M)^2)$ and the radiative recoil $O(\alpha m/M)$, is [9, 22, 31, 32]

$$g_{\text{rec}} = \frac{m}{M} \frac{(Z\alpha)^2}{4} \left[1 + \frac{11}{48} (Z\alpha)^2 - \frac{m}{M} (1+Z) - \frac{\alpha}{3\pi} \right]. \quad (\text{A8})$$

The finite nuclear size correction including the corresponding QED contribution is

$$g_N = \frac{2}{5} (Z\alpha R_{\text{sph}})^{2\gamma} \frac{(Z\alpha)^2}{2} \left[1 + (Z\alpha)^2 H_N^{(0,2+)} \right] \times \left[1 + \frac{\alpha}{\pi} G_{\text{NQED}} \right], \quad (\text{A9})$$

where $\gamma = \sqrt{1 - (Z\alpha)^2}$ and $R_{\text{sph}} = \sqrt{5/3} R$ the radius of the nuclear sphere with the root-mean-square radius R , and the remainder functions $H_N^{(0,2+)}$ and G_{NQED} were evaluated in Refs. [7, 21, 33].

Appendix B: Fitting of the $1/Z$ expansion coefficients

In this section we describe the fitting procedure used for the identification of the coefficients of the $1/Z$ expansion.

The general task is to fit a data set of n points (Z_i, F_i) , $i = 1, \dots, n$, to the following model function with N ($N < n$) parameters,

$$f_N(Z) = \sum_{k=0}^{N-1} c_k Z^{a-k}, \quad (\text{B1})$$

where a is the exponent of the leading term of the $1/Z$ expansion.

In order to find the optimal values of the fitting parameters c_k , we use the weighted least-squares regression. Specifically, we minimize the functional

$$S_N = \sum_{i=1}^n \frac{(F_i - f_N(Z_i))^2}{\delta F_i^2 + \sigma_N^2(Z_i)}, \quad (\text{B2})$$

where δF_i are the numerical errors of F_i and $\sigma_N(Z)$ is the estimate of the error due to the truncation of the $1/Z$ expansion in the fitting function, taken as the last term of the fitting ansatz divided by Z ,

$$\sigma_N(Z) = c_{N-1} Z^{a-N}. \quad (\text{B3})$$

In practice, we make our fit in two steps. First, we perform the least-square regression without weights. The obtained value of the c_{N-1} coefficient is then used for the estimation of the truncation error in the weighted least-square regression performed on the second step.

In the cases relevant for the present work, one or two first coefficients of the $1/Z$ expansion are known analytically. We use this fact in order to access the errors of our fitting procedure. First, we treat one of the known coefficients as a free fitting parameter and select three different fitting functions that give the best approximation to the known result. After that, we set the known coefficients to their exact values, perform the fit with the three fitting functions, and finally take the average of the three results and the maximal deviation between them as the final value and its error, respectively.

Fitting the results for the α^2 and α^3 corrections (whose numerical accuracy is very high), we used fitting functions with 8-10 parameters. For the recoil corrections (whose accuracy is much lower), we used 5-6 fitting parameters.

-
- [1] S. Sturm, A. Wagner, B. Schabinger, J. Zatorski, Z. Harman, W. Quint, G. Werth, C. H. Keitel, and K. Blaum, Phys. Rev. Lett. **107**, 023002 (2011).
 - [2] S. Sturm, A. Wagner, M. Kretzschmar, W. Quint, G. Werth, and K. Blaum, Phys. Rev. A **87**, 030501 (2013).
 - [3] S. Sturm, F. Köhler, J. Zatorski, A. Wagner, Z. Harman, G. Werth, W. Quint, C. H. Keitel, and K. Blaum, Nature **506**, 467470 (2014).
 - [4] P. J. Mohr, D. B. Newell, and B. N. Taylor, Rev. Mod. Phys. **88**, 035009 (2016).
 - [5] A. Wagner, S. Sturm, F. Köhler, D. A. Glazov, A. V. Volotka, G. Plunien, W. Quint, G. Werth, V. M. Shabaev, and K. Blaum, Phys. Rev. Lett. **110**, 033003 (2013).
 - [6] F. Köhler et al., Nature communications **7** (2016).
 - [7] V. A. Yerokhin, E. Berseneva, Z. Harman, I. I. Tupitsyn, and C. H. Keitel, Phys. Rev. Lett. **116**, 100801 (2016).
 - [8] V. M. Shabaev, D. A. Glazov, M. B. Shabaeva, V. A. Yerokhin, G. Plunien, and G. Soff, Phys. Rev. A **65**, 062104 (2002).
 - [9] D. A. Glazov, V. M. Shabaev, I. I. Tupitsyn, A. V. Volotka, V. A. Yerokhin, G. Plunien, and G. Soff, Phys. Rev. A **70**, 062104 (2004).
 - [10] D. A. Glazov, A. V. Volotka, V. M. Shabaev, I. I. Tupitsyn, and G. Plunien, Phys. Lett. A **357**, 330 (2006).
 - [11] D. A. Glazov, A. V. Volotka, V. M. Shabaev, I. I. Tupitsyn, and G. Plunien, Phys. Rev. A **81**, 062112 (2010).
 - [12] A. V. Volotka, D. A. Glazov, V. M. Shabaev, I. I. Tupitsyn, and G. Plunien, Phys. Rev. Lett. **112**, 253004 (2014).
 - [13] R. A. Hegstrom, Phys. Rev. A **7**, 451 (1973).
 - [14] Z.-C. Yan, Phys. Rev. Lett. **86**, 5683 (2001).
 - [15] Z.-C. Yan, J. Phys. B **35**, 1885 (2002).
 - [16] M. Puchalski and K. Pachucki, Phys. Rev. A **73**, 022503 (2006).

TABLE III. Binding corrections to the g -factor of the ground state of $^{28}\text{Si}^{11+}$. “LO” denotes the lowest-order (in $Z\alpha$) contribution of the corresponding correction, to be multiplied by the prefactor specified in the second column. “HO” denotes the higher-order (in $Z\alpha$) contribution, to be multiplied by the prefactor specified in the second column. δg denotes the contribution to the g -factor, obtained as the sum of the lowest-order and the higher-order contributions, multiplied by the prefactor. Nuclear parameters used in the calculation are: $M/m = 50\,984.832\,73$ and $R = 3.1224(24)$ fm.

Order	Prefactor	LO	HO	$\delta g \times 10^6$
Electron-electron interaction:				
$1/Z^0$	$\alpha^2 Z^2$	-0.166 666 667	-0.000 546 606	-1 745.249 323
$1/Z^1$	$\alpha^2 Z$	0.429 812 529	0.001 552 492	321.590 803
$1/Z^2$	α^2	-0.128 204 (9)	-0.000 92 (1)	-6.876 0 (5)
$1/Z^{3+}$	$\alpha^2 Z^{-1}$	0.024 8 (1)	0.000 0 (3)	0.094 (1)
One-loop QED:				
$1/Z^0$	$\alpha^3 Z^2$	0.013 262 912	0.002 813 49 (4)	1.224 449 (3)
$1/Z^1$	$\alpha^3 Z$	-0.039 879 702	-0.005 1 (9)	-0.245 (5)
$1/Z^{2+}$	α^3	0.022 467 862	0.000 (4)	0.009 (2)
Recoil:				
$1/Z^0$	$\alpha^2 (m/M) Z^2$	0.250 000 000	0.001 (8)	0.051 (2)
$1/Z^{1+}$	$\alpha^2 (m/M) Z$	-0.832 9 (1)	0.00 (4)	-0.012 2 (6)
Two-loop QED:				
$1/Z^0$	$\alpha^4 Z^2$	-0.002 773 485	-0.000 6 (6)	-0.001 9 (3)
$1/Z^1$	$\alpha^4 Z$	0.008 339 479	0.000 (3)	0.000 3 (1)
Finite nuclear size:				
$1/Z^0$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z^2$	0.200 000 000	0.001 90 (7)	0.002 574 (4)
$1/Z^1$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z$	-0.570 2 (2)	-0.008 0 (3)	-0.000 527 (1)
$1/Z^2$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2$	0.214 (5)	0.002 (3)	0.000 014
Radiative recoil:				
$1/Z^0$	$\alpha^3 (m/M) Z^2$	-0.026 525 824	0.000 (1)	-0.000 040 (2)
$1/Z^{1+}$	$\alpha^3 (m/M) Z$	0.040 337 (6)	0.000 (2)	0.000 004
Second-order recoil:				
$1/Z^0$	$\alpha^2 (m/M)^2 Z^2$	-3.750 000 000	0.0 (2)	-0.000 015 (1)
≥ 3 -loop QED:				
$1/Z^0$	$\alpha^5 Z^2$	0.0031 629 03	0.000 0 (7)	0.000 013 (3)
Total $g - g_e$:				
Theory, this work				-1 429.412 (6)
Theory [12]				-1 429.412 (8)
Experiment [5]				-1 429.414 5 (21)

- [17] M. Puchalski and K. Pachucki, Phys. Rev. A **78**, 052511 (2008).
- [18] M. Puchalski and K. Pachucki, Phys. Rev. A **92**, 012513 (2015).
- [19] D. Hanneke, S. Fogwell, and G. Gabrielse, Phys. Rev. Lett. **100**, 120801 (2008).
- [20] V. M. Shabaev and V. A. Yerokhin, Phys. Rev. Lett. **88**, 091801 (2002).
- [21] V. A. Yerokhin, E. Berseneva, Z. Harman, I. I. Tupitsyn, and C. H. Keitel, Phys. Rev. A **94**, 022502 (2016).
- [22] H. Grotch and R. A. Hegstrom, Phys. Rev. A **4**, 59 (1971), [(E) Phys. Rev. A **8**, 575 (1973)].
- [23] K. Pachucki, U. D. Jentschura, and V. A. Yerokhin, Phys. Rev. Lett. **93**, 150401 (2004), [erratum: *ibid.*, **94**, 229902 (2005)].
- [24] K. Pachucki, A. Czarnecki, U. D. Jentschura, and V. A. Yerokhin, Phys. Rev. A **72**, 022108 (2005).
- [25] V. A. Yerokhin and Z. Harman, <http://arxiv.org/abs/1704.08080>.
- [26] S. G. Karshenboim, R. N. Lee, and A. I. Milstein, Phys. Rev. A **72**, 042101 (2005).
- [27] R. N. Lee, A. I. Milstein, I. S. Terekhov, and S. G. Karshenboim, Phys. Rev. A **71**, 052501 (2005).
- [28] A. Czarnecki and R. Szafron, Phys. Rev. A **94**, 060501 (2016).
- [29] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys. Rev. D **91**, 033006 (2015).
- [30] S. Laporta, <https://arxiv.org/abs/1704.06996>.
- [31] R. Faustov, Phys. Lett. **33B**, 422 (1970).
- [32] F. E. Close and H. Osborn, Phys. Lett. **34B**, 400 (1971).
- [33] V. A. Yerokhin, C. H. Keitel, and Z. Harman, J. Phys. B **46**, 245002 (2013).

TABLE IV. Individual binding correction to the g -factor of the ground state of $^{16}\text{O}^{5+}$. Notations are the same as in Table III. Nuclear parameters used in the calculation are: $M/m = 29\,148.949\,75$ and $R = 2.6991(52)$ fm.

Order	Prefactor	LO	HO	$\delta g \times 10^6$
Electron-electron interaction:				
$1/Z^0$	$\alpha^2 Z^2$	-0.166 666 667	-0.000 177 823	-568.620 484
$1/Z^1$	$\alpha^2 Z$	0.429 812 529	0.000 505 635	183.320 201
$1/Z^2$	α^2	-0.128 204 (9)	-0.000 3 (2)	-6.843 (8)
$1/Z^{3+}$	$\alpha^2 Z^{-1}$	0.020 69 (7)	0.000 00 (7)	0.137 7 (7)
One-loop QED:				
$1/Z^0$	$\alpha^3 Z^2$	0.013 262 912	0.001 330 41 (3)	0.362 936 (1)
$1/Z^1$	$\alpha^3 Z$	-0.039 879 702	-0.002 (2)	-0.129 (5)
$1/Z^{2+}$	α^3	0.022 539 315	0.000 (1)	0.008 8 (5)
Recoil:				
$1/Z^0$	$\alpha^2 (m/M) Z^2$	0.250 000 000	0.000 (2)	0.029 3 (2)
$1/Z^{1+}$	$\alpha^2 (m/M) Z$	-0.811 399 (3)	0.000 (9)	-0.011 9 (1)
Two-loop QED:				
$1/Z^0$	$\alpha^4 Z^2$	-0.002 773 485	-0.000 05 (6)	-0.000 51 (1)
$1/Z^1$	$\alpha^4 Z$	0.008 339 479	0.000 0 (2)	0.000 189 (5)
Finite nuclear size:				
$1/Z^0$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z^2$	0.200 000 000	0.000 42 (6)	0.000 194 (1)
$1/Z^1$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z$	-0.570 2 (2)	-0.002 4 (3)	-0.000 069
$1/Z^2$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2$	0.214 (5)	0.001 (1)	0.000 003
Radiative recoil:				
$1/Z^0$	$\alpha^3 (m/M) Z^2$	-0.026 525 824	0.000 0 (3)	-0.000 023
$1/Z^{1+}$	$\alpha^3 (m/M) Z$	0.040 504 18 (6)	0.000 0 (5)	0.000 004
Second-order recoil:				
$1/Z^0$	$\alpha^2 (m/M)^2 Z^2$	-2.250 000 000	0.00 (3)	-0.000 009
≥ 3 -loop QED:				
$1/Z^0$	$\alpha^5 Z^2$	0.003 162 903	0.000 00 (4)	0.000 004
Total $g - g_e$:				
Theory, this work				-391.745 9 (96)
Theory [9]				-391.700 (32)

TABLE V. Individual binding correction to the g -factor of the ground state of $^{12}\text{C}^{3+}$. Notations are the same as in Table III. Nuclear parameters used in the calculation are: $M/m = 21\,868.663\,86$ and $R = 2.4702(24)$ fm.

Order	Prefactor	LO	HO	$\delta g \times 10^6$
Electron-electron interaction:				
$1/Z^0$	$\alpha^2 Z^2$	-0.166 666 667	-0.000 099 947	-319.699 730
$1/Z^1$	$\alpha^2 Z$	0.429 812 529	0.000 284 265	137.419 421
$1/Z^2$	α^2	-0.128 204 (9)	-0.000 17 (8)	-6.836 (4)
$1/Z^{3+}$	$\alpha^2 Z^{-1}$	0.016 65 (5)	0.000 00 (3)	0.147 8 (6)
One-loop QED:				
$1/Z^0$	$\alpha^3 Z^2$	0.013 262 912	0.000 879 11 (3)	0.197 838
$1/Z^1$	$\alpha^3 Z$	-0.039 879 702	-0.000 9 (9)	-0.095 (2)
$1/Z^{2+}$	α^3	0.022 618 936	0.000 0 (8)	0.008 8 (3)
Recoil:				
$1/Z^0$	$\alpha^2 (m/M) Z^2$	0.250 000 000	0.000 1 (9)	0.021 93 (8)
$1/Z^{1+}$	$\alpha^2 (m/M) Z$	-0.793 800 (1)	0.000 (4)	-0.011 60 (6)
Two-loop QED:				
$1/Z^0$	$\alpha^4 Z^2$	-0.002 773 485	0.000 02 (2)	-0.000 281(2)
$1/Z^1$	$\alpha^4 Z$	0.008 339 479	0.000 00 (8)	0.000 142 (1)
Finite nuclear size:				
$1/Z^0$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z^2$	0.200 000 000	0.000 18 (9)	0.000 051
$1/Z^1$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2 Z$	-0.570 2 (2)	-0.001 4 (7)	-0.000 024
$1/Z^2$	$(Z\alpha R_{\text{sph}})^{2\gamma} \alpha^2$	0.214 (5)	0.000 4 (8)	0.000 002
Radiative recoil:				
$1/Z^0$	$\alpha^3 (m/M) Z^2$	-0.026 525 824	0.000 0 (1)	-0.000 017
$1/Z^{1+}$	$\alpha^3 (m/M) Z$	0.040 698 000	0.000 0 (2)	0.000 004
Second-order recoil:				
$1/Z^0$	$\alpha^2 (m/M)^2 Z^2$	-1.750000000	0.00 (1)	-0.000007
≥ 3 -loop QED:				
$1/Z^0$	$\alpha^5 Z^2$	0.003 162 903	0.000 0 (1)	0.000 002
Total $g - g_e$:				
Theory, this work				-188.847 (5)
Theory [9]				-188.819 (19)